

Association between a Vasopressin Receptor AVPR1A Promoter Region Microsatellite and Eating Behavior Measured by a Self-Report Questionnaire (Eating Attitudes Test) in a Family-Based Study of a Nonclinical Population

Rachel Bachner-Melman,^{1,2} Ada H. Zohar,³ Yoel Elizur,¹ Lubov Nemanov,⁴
Inga Gritsenko,⁴ Daphnie Konis,² and Richard P. Ebstein^{2,4*}

¹Department of Psychology, Mount Scopus, Hebrew University, Jerusalem, Israel

²Scheinfeld Center of Human Genetics for the Social Sciences, Department of Psychology,
Hebrew University, Jerusalem, Israel

³Department of Psychology and Behavioral Sciences, Ruppin Academic Center,
Emek Hefer, Israel

⁴Sarah Herzog Memorial Hospital, Jerusalem, Israel

Accepted 11 January 2004

Abstract: Objectives: Considerable evidence including twin and family studies suggests that biologic determinants interact with cultural cues in the etiology of anorexia and bulimia nervosa. A gene that makes “biologic sense” in contributing susceptibility to these disorders, and to our knowledge not previously investigated for this phenotype, is the vasopressin receptor (AVPR1A), which we have tested for association with eating pathology. **Methods:** We genotyped 280 families with same-sex siblings for two microsatellites in the promoter region of the AVPR1A gene. Siblings completed the 26-item Eating Attitudes Test (EAT) and the Drive for Thinness (DT) and Body Dissatisfaction (BD) subscales of the Eating Disorders Inventory (EDI). The Quantitative Transmission Disequilibrium Test program (QTDT), which employs flexible and powerful variance-components procedures, was used to test for an association between EAT scores and the two AVPR1A promoter region microsatellites, RS1 and RS3. **Results:** A significant association ($p = .036$) was detected between the RS3 microsatellite and EAT scores. The strongest association was between RS3 and the Dieting subscale of the EAT ($p = .011$). A significant association was also observed between the EDI-DT and the RS3 microsatellite ($p = .0450$). **Conclusions:** We demonstrate for the first time an association between a microsatellite polymorphism in the AVPR1A promoter region and

Supported, in part, by the Israel Science Foundation founded by the Israel Academy of Sciences and Humanities (RPE), the Israel Association of University Women (RB-M), and the Levin Center for the Normal and Psychopathological Development of the Child and Adolescent (RB-M).

*Correspondence to: Dr. Richard P. Ebstein, Scheinfeld Center of Human Genetics for the Social Sciences, Department of Psychology, Hebrew University, P.O. Box 35300, Jerusalem 91351, Israel.
E-mail: ebstein@mscc.huji.ac.il

Published online in Wiley InterScience (www.interscience.wiley.com). DOI: 10.1002/eat.20049

scores on the EAT as well as with the EDI-DT. The strongest association was observed between the RS3 microsatellite and the Dieting subscale of the EAT. The relevant phenotype appears to tap severe dietary restriction for weight loss purposes. © 2004 by Wiley Periodicals, Inc. *Int J Eat Disord* 36: 451–460, 2004.

Key words: AVPR1A; Eating Attitudes Test; twin and family studies; microsatellite polymorphisms

INTRODUCTION

Anorexia nervosa (AN) and bulimia nervosa (BN) are disorders characterized by abnormal patterns of eating behavior and by disturbances in attitudes and perceptions toward body weight and shape (Kaye, Klump, Frank, & Strober, 2000). In Western societies, sociocultural factors such as thinness norms clearly play a crucial role in shaping eating pathology (Botta, 1999; Garner & Garfinkel, 1980; Stice, Schupak-Neuberg, Shaw, & Stein, 1994). However, the low prevalence of severe eating disorders, a marked gender bias, and early age of onset suggest that other factors, namely, biologic determinants, interact with cultural cues in the etiology of AN and BN. Twin and family studies have suggested significant heritability for these disorders (Bulik, Sullivan, Wade, & Kendler, 2000; Holland, Sicotte, & Treasure, 1988; Klump, Miller, Keel, McGue, & Iacono, 2001; Walters & Kendler, 1995).

Both candidate gene (Hinney, Remschmidt, & Hebebrand, 2000; Tozzi, Bergen, & Bulik, 2002) and linkage strategies (Devlin et al., 2002; Grice et al., 2002) are currently being applied in an effort to identify genes and chromosomal regions contributing susceptibility to AN and BN, with promising preliminary results. Recently, a Japanese group used an innovative strategy in an attempt to identify genes contributing to AN and BN (Matsushita, Nakamura, Nishiguchi, & Higuchi, 2002). They found provisional evidence for an association between the serotonin transporter promoter region polymorphism (Lesch et al., 1996) and scores on the 26-item Eating Attitudes Test (EAT) in a group of unrelated female nursing students. Eating pathology in this scheme is operationalized by a score above 20 on the EAT, which is widely accepted as a reliable and valid self-report screening instrument for eating disorders (Garner, 1991; Garner & Garfinkel, 1979; Garner, Olmsted, Bohr, & Garfinkel, 1982). The clinical success of the EAT as a screening tool suggests that eating disorders may be the extremes of a behavioral spectrum, just as certain ranges in normal personality traits may indicate personality disorders (Livesley, Jang, & Vernon, 1998). Scores on the EAT show substantial heritability, with approximately one-half the variance between individuals attributable to genes (Rutherford, McGuffin, Katz, & Murray, 1993). It seems biologically meaningful to surmise that some of the quantitative trait loci partially determining EAT scores may contribute to AN and BN.

A gene of potential interest in contributing susceptibility to eating disorders, and to our knowledge not previously investigated for this phenotype, is the vasopressin receptor (AVPR1A). Arginine vasopressin (AVP), the endogenous ligand for the V1a vasopressin receptor, is synthesized in a number of sites in the brain, including the paraventricular, supraoptic, and suprachiasmatic nuclei of the hypothalamus (Barberis & Tribollet, 1996; Tribollet, Arsenijevic, & Barberis, 1998). Vasopressin is known for its anorectic effect in animal studies (Langhans, Delprete, & Scharrer, 1991). In the brain, AVP participates in glucose homeostasis as a neurotransmitter or neuromodulator, exciting neurons in the

paraventricular nucleus (Inenaga, Dyball, Okuya, & Yamashita, 1986; Inenaga & Yamashita, 1986) and ventromedial glucose-responsive neurons in the hypothalamus (Kow & Pfaff, 1986). Direct administration of micro doses of AVP into the nucleus of the tractus solitarius of anesthetized or awake rats rapidly increased the levels of blood glucose concentration and brain arteriovenous glucose difference (Yarkov, Montero, Lemus, Roces de Alvarez-Buylla, & Alvarez-Buylla, 2001). AVP also appears to mediate the effects of several neuropeptides on feeding behavior (Backberg, Hervieu, Wilson, & Meister, 2002; Dhillon et al., 2003; Niimi, Murao, & Taminato, 2001; Ur, Wilkinson, Morash, & Wilkinson, 2002; Wren et al., 2002).

In addition, a number of human studies suggest that vasopressin may play a role in the pathophysiology of AN and BN (Kaye, 1996). Elevated levels of AVP were observed in the cerebrospinal fluid of women recovered from AN and BN (Frank, Kaye, Altemus, & Greeno, 2000). Low serum activity of prolyl endopeptidase, a cytosolic endopeptidase that cleaves peptide bonds on the carboxyl side of proline in proteins such as vasopressin, has been observed in AN (Maes et al., 2001).

In the current study, we genotyped 280 families with same-sex siblings for two microsatellites (Thibonnier et al., 2000) in the promoter region of the *AVPR1A* gene. Scores on the EAT and the Drive for Thinness (DT) and Body Dissatisfaction (BD) subscales of the Eating Disorders Inventory-2 (EDI) were available for these subjects. The Quantitative Transmission Disequilibrium Test program (QTDT; Abecasis, Cardon, & Cookson, 2000; Abecasis, Cookson, & Cardon, 2000), which employs flexible and powerful variance-components procedures, was used to test for an association between EAT scores and the two *AVPR1A* promoter region microsatellites.

METHODS

Subjects

Respondents were primarily college students at Israeli colleges and their families, recruited by word of mouth and advertisements on campus noticeboards. The sample analyzed in the current study included 280 families, each with two parents and same-sex siblings between the ages of 14 and 34 (average age = 22.73 ± 4.09). Pedigrees include 7 single-sibling families, 213 two-sibling families, 41 families with three siblings, 12 families with four siblings, 6 families with five siblings, and 1 family with six siblings.

Each contact person received a number of questionnaires (equal to the number of participating siblings) and two sterile test tubes per family member for DNA sampling, containing 10 ml of Aquafresh mouthwash. Questionnaires were completed by siblings but not parents. After a complete description of the study to the subjects, written informed consent was obtained. Completed questionnaires and DNA mouthwashes were returned by mail or hand delivered to an office. The contact person received a modest monetary incentive and the study was approved by the local institutional review board and by the genetics committee of the Israeli Ministry of Health.

Instruments

EAT

The EAT is a 26-item self-report factor-analytically derived scale, originally validated on a sample of 160 women with eating disorders and 140 female nonclinical controls (Garner et al., 1982). It is reliable and valid, correlates highly with the original 40-item

scale ($r = .98$; Garner et al., 1982), and screens for cases of eating disorders in both clinical and nonclinical populations. The EAT is scored on a 6-point Likert scale with answers ranging from *never* to *always*.

EDI

The EDI-2 is a self-report measure of symptoms generally related to eating disorders (Garner, 1991). It contains 11 standardized subscales, each independently derived and representing a unique trait. In the current study, the EDI was used to assess participants' body dissatisfaction and drive for thinness. These subscales have excellent internal consistency, content validity, criterion-based and construct validity, and good test-retest reliability in eating-disordered and healthy control subjects (Garner, 1991).

DNA Extraction and Genotyping

DNA samples were obtained from all family members and extracted using the Master Pure kit (Epicentre, Madison WI). Amplification of the RS1 and RS3 vasopressin was achieved using the following pair of primers: RS1 (corresponds to the GATA microsatellite at position -553 relative to the start site as first described by Thibonnier et al. (2000): forward (fluorescent) 5' AGG GAC TGG TTC TAC AAT CTG C 3'; reverse 5' ACC TCT CAA GTT ATG TTG GTG G 3'. RS3 (corresponds to the GT dinucleotide repeat at -3956 (Thibonnier et al., 2000); forward (fluorescent) 5' CCT GTA GAG ATG TAA GTG CT 3'; reverse 5' TCT GGA AGA GAC TTA GAT GG 3'. The two microsatellites were somewhat in linkage disequilibrium ($D' = .224$). D' was calculated using the method of Slatkin and Excoffier (1996), which can be downloaded from the following internet site (Abecasis & Cookson, 2000): <http://www.sph.umich.edu/csg/abecasis/GOLD/index.html>.

Each reaction mixture contained 0.5 μ M primer. A ReddyMix master mix was used (Abgene, Surrey, UK) at a magnesium concentration of 1.5–2.5 mM $MgCl_2$. The sample was initially heated at 95°C for 5' followed by 30 cycles of 95°C (30 sec), 55°C (30 sec), 72°C (40 sec), and a final extension step of 72°C for 10 min. The polymerase chain reaction product was analyzed on an ABI 310 DNA analyzer (Applied Biosystems, Foster City, CA).

Statistical Analysis

QTDT is used to analyze quantitative or discrete traits in nuclear families, with or without parental genotypes, or in extended pedigrees. The tests described by Allison (1997), Rabinowitz (1997), Monks, Kaplan, and Weir (1998), and Fulker, Cherny, Sham, and Hewitt (1999) are also supported. Because simple models of association do not provide valid tests of linkage disequilibrium when multiple offspring per family are considered, we used the robust variance-components procedures as detailed in the QTDT package. We examined an association between two microsatellites in the promoter region of the *AVPR1A* gene (RS1 and RS3) and scores on the EAT and the DT and BD subscales of the EDI. Age and body mass index (BMI) were entered as covariates. Because entering gender as a covariate had no effect on results, gender is not shown as a covariate in Table 1. No correction was made for multiple comparisons (two microsatellites examined \times two eating scales) and the results should therefore be interpreted cautiously.

We carried out a power analysis using PBAT (Lange & Laird, 2002). The following parameters were entered in the model: continuous trait, additive genetic model, allele frequency of the disease gene = .2, heritability = .05, significance level = .05, allele frequency of the marker gene = .15, P (disease allele A/marker allele A) = .9, $D' = .875$, 280 families with two siblings. Power was calculated to be .754.

RESULTS

Table 1 shows the global *p* value (corrected for multiple alleles of the microsatellite) when the *AVPR1A* microsatellites were tested for an association with the EAT and EDI scores. A significant association (*p* = .036) was detected between the RS3 microsatellite and EAT scores. Therefore, we examined the association for each of the EAT subscales. The strongest association was observed between RS3 and the Dieting subscale (*p* = .011). Because the EAT Dieting subscale and the EDI-DT subscale correlate very highly, we also examined the EDI-DT for association with the *AVPR1A* receptor polymorphism. As expected, a significant association was observed between the EDI-DT and the RS3 (*p* = .0450).

Values for individual alleles and the association with EAT scores are shown in Table 2. The allele frequencies of the RS1 and RS3 microsatellites are similar to what has previously been reported (Thibonnier et al., 2000).

Table 1. Testing for association between EAT scores and *AVPR1A* microsatellite polymorphisms

Testing trait		EAT				
Testing marker		RS1				
Allele	df(0)	LnLk(0)	df(T)	LnLk(T)	ChiSq	<i>p</i>
All	609	2659.62	604	2656.86	5.51	
Testing marker		RS3				
Allele	df(0)	LnLk(0)	df(T)	LnLk(T)	ChiSq	<i>p</i>
All	611	2668.15	606	2662.21	11.87	.036
Testing trait		Dieting subscale				
Testing marker		RS1				
Allele	df(0)	LnLk(0)	df(T)	LnLk(T)	ChiSq	<i>p</i>
All	609	2436.50	604	2434.05	4.91	
Testing marker		RS3				
Allele	df(0)	LnLk(0)	df(T)	LnLk(T)	ChiSq	<i>p</i>
All	611	2447.30	606	2439.92	14.76	.0114
Testing trait		EDI Drive for thinness				
Testing marker		RS1				
Allele	df(0)	LnLk(0)	df(T)	LnLk(T)	ChiSq	<i>p</i>
All	608	2190.01	603	2186.71	6.61	
Testing marker		RS3				
Allele	df(0)	LnLk(0)	df(T)	LnLk(T)	ChiSq	<i>p</i>
All	610	2201.27	605	2195.62	11.30	.0458

Note: The null and full models are evaluated. The null model: means = $\mu + \text{covariate_age} + \text{covariate_BMI} + B$; variances = $V_e + V_g + V_a$. The full model: means = $\mu + \text{covariate_age} + \text{covariate_BMI} + B + W$; variances = $V_e + V_g + V_a$.

B is a between-family component of association, whereas W is a within-family component of association. Alleles with rare frequencies of 0.05% or less are lumped together. The individual effects for all other alleles are estimated producing a single, global *p* value. A fuller explanation of these models can be found at <http://www.sph.umich.edu/csg/abecasis/> as well as in the cited references. In the model, e = nonshared environment; g = polygenic effects—a function of relatedness between family members and may be due to polygenes; a = additive genetic effects. The chi-square value for determining the overall global *p* value is derived by subtracting the chi-square value of the null model from the chi-square value of the full model. EAT = Eating Attitudes Test.

Table 2. Testing for association between EAT scores and *AVPR1A* microsatellite polymorphisms individual alleles

Testing marker		RS1					
Allele	df(0)	LnLk(0)	df(T)	LnLk(T)	ChiSq	<i>p</i>	allele frequency
1	613	2662.75	612	2662.15	1.19		0.028
2	613	2661.75	612	2660.98	1.54		0.144
3	613	2662.22	612	2662.16	0.13		0.299
4	613	2662.04	612	2661.83	0.43		0.277
5	613	2661.18	612	2661.10	0.15		0.128
6	613	2662.72	612	2660.55	4.35	.0369	0.094
7	*** Not tested ***						0.019
8	*** Not tested ***						0.006
9	*** Not tested ***						0.004
Testing marker		RS3					
Allele	df(0)	LnLk(0)	df(T)	LnLk(T)	ChiSq	<i>p</i>	allele frequency
1	*** Not tested ***						0.006
2	615	2669.09	614	2669.05	0.08		0.043
3	615	2669.12	614	2667.02	4.21	.0403	0.131
4	615	2669.04	614	2668.27	1.54		0.244
5	615	2669.09	614	2666.10	5.99	.0144	0.192
6	615	2669.05	614	2668.12	1.86		0.136
7	615	2668.75	614	2668.05	1.40		0.098
8	615	2669.09	614	2664.66	8.86	.0029	0.024
9	615	2668.61	614	2668.31	0.60		0.024
10	615	2669.13	614	2667.66	2.94	.0864	0.061
11	*** Not tested ***						0.014
12	*** Not tested ***						0.002
13	*** Not tested ***						0.001
14	*** Not tested ***						0.008
15	*** Not tested ***						0.003
16	*** Not tested ***						0.009

Note: EAT = Eating Attitudes Test.

DISCUSSION

The regulation of food intake has been shown to be partially mediated by a variety of neuroregulators including neuropeptide Y, melanocortin, orexins, opioid peptides, leptins, ghrelin, cholecystokinin, and AVP (Beck, 2000; Saper, Chou, & Elmquist, 2002). In the current study, we demonstrate an association between a microsatellite polymorphism in the *AVPR1A* promoter region and scores on the EAT as well as with the EDI-DT, in respondents recruited from college campuses and the community. Significant, albeit modest, associations were observed using a robust family-based design in a variance-components framework suitable for family pedigrees of various sizes (Abecasis, Cardon, et al., 2000; Abecasis, Cookson, et al., 2000). The strongest association was observed between the *AVPR1A* promoter region microsatellite RS3 and the Dieting subscale of the EAT. No significant association was observed between the RS1 microsatellite and any of the scales administered or between the RS3 microsatellite and the Bulimia and Oral Control subscales of the EAT or the BD subscale of the EDI.

Whereas high scores on the Bulimia subscale of the EAT are indicative of bulimic but not restrictive eating disorders, high scores on the Dieting subscale are a feature of both (Garner, Olmsted, & Garfinkel, 1985). Given the high correlation between the EAT Dieting subscale and the EDI-DT ($r = .912$), it is not surprising that an association (weaker but,

nevertheless, significant) was observed between the RS3 microsatellite and both the Dieting subscale of the EAT and the EDI-DT. The behavioral phenotype in question therefore appears to be one of dietary restriction which, taken to an extreme, characterizes AN.

Several mechanisms related to the physiology of vasopressin action in the brain and its role in the regulation of food intake might explain the observed association. First, vasopressin appears to be involved in the regulation of glucose homeostasis (Kow & Pfaff, 1986), suggesting that it may be involved in the "energy deficit" initiation of food intake. Second, vasopressin may affect dieting behavior via its action on the reward properties of food or lack of food, which is, in turn, closely related to glucose regulation. Under certain conditions, for example, vasopressin and related peptides decrease brain reward and addictive behavior in experimental studies in animal and humans (Van Ree, 1986). Of relevance to this suggestion is the finding that eating-disordered patients, especially patients with AN, often report subjective transient symptomatic improvement or even feelings of euphoria as a result of dieting and food restriction. The food restriction is consequently reinforced by these feelings (Kaye & Strober, 1999).

Third, the *AVPR1A* receptor may contribute to eating behavior via the mechanism of stress-induced feeding problems, because vasopressin is involved in the leptin-induced activation of the hypothalamic pituitary axis (HPA) axis (Levine, 2001; Morimoto et al., 2000). The possibility of a stress axis factor is supported by publications suggesting the presence of pre-onset stressors and coping difficulties in the etiology of AN (Horesh et al., 1996; Welch, Upchurch, & Loscalzo, 1997).

Most of the information regarding the contribution of *AVPR1A* to social behavior is based on studies of the vole (Wang, Young, De Vries, & Insel, 1998). The prairie vole is highly social and monogamous, in striking contrast to the asocial and no-monogamous montane vole. There is evidence that microsatellite instability in the promoter region of this and other genes may be a major contributing factor to diversity in both region-specific gene expression and resulting behavioral phenotypes (Hammock & Young, 2002). In humans, the function of the microsatellites in regulating the *AVPR1A* gene in humans is still poorly understood. However, Hammock and Young (2002) suggested that "minor variations in microsatellite length at the *AVPR1A* locus in humans can contribute to the diversity in human behavioral traits that are considered to be in the 'normal' range of behavior" (p 401). Recently, the promoter region microsatellites have been linked to autism, providing preliminary evidence that the *AVPR1A* gene may play a role in social communication and behavior in humans (Kim et al., 2002).

Vasopressin also profoundly affects social interactions, social communication, and pair-bonding in a variety of vertebrate species (Insel & Young, 2000). Because there is ample evidence that eating style is influenced by social context across cultural groups (Backman, Haddad, Lee, Johnston, & Hodgkin, 2002; Clendenen, Herman, & Polivy, 1994; Monge-Rojas, Nunez, Garita, & Chen-Mok, 2002; Satia-Abouta, Patterson, Kristal, Teh, & Tu, 2002), vasopressin and its receptor could also affect human eating behavior via its action as a "social" hormone. It is noteworthy that in *Caenorhabditis elegans*, a neuropeptide receptor, albeit a different one, controls the choice between solitary and social feeding behavior patterns (Coates & de Bono, 2002).

Eating habits in animals and humans reflect many factors beyond simple energy requirements needed for the maintenance and growth of the organism (Levine & Billington, 1997). When, why, and how much people eat is affected by food availability and palatability, boredom, emotional state, environmental stimuli (Birch, McPhee, Sullivan, & Johnson, 1989), and the presence of others (de Castro & de Castro, 1989). Eating behavior is simultaneously primitive and sophisticated. It is inextricably

entrenched in social settings and rituals, in culinary and consumption customs, and in cultural habits and norms. Whereas in contemporary Western cultures dieting and the pursuit of thinness are desirable, fashionable, and widespread, they are also clearly predictive of the development of eating disorders (Garner & Garfinkel, 1980; Hsu, 1990; Polivy & Herman, 1987). Our results should meanwhile be interpreted cautiously and regarded as preliminary. Future research with independent samples should seek to replicate the evidence we present for the involvement of vasopressin and the *AVPR1A* receptor in feeding behavior, particularly in the socially approved, yet often maladaptive, practice of dieting and food restriction.

REFERENCES

- Abecasis, G.R., Cardon, L.R., & Cookson, W.O. (2000). A general test of association for quantitative traits in nuclear families. *American Journal of Human Genetics*, 66(1), 279–292.
- Abecasis, G.R., & Cookson, W.O. (2000). GOLD—graphical overview of linkage disequilibrium. *Bioinformatics*, 16(2), 182–183.
- Abecasis, G.R., Cookson, W.O., & Cardon, L.R. (2000). Pedigree tests of transmission disequilibrium. *European Journal of Human Genetics*, 8(7), 545–551.
- Allison, D.B. (1997). Transmission-disequilibrium tests for quantitative traits [published erratum appears in *Am J Hum Genet* 1997 Jun;60(6):1571]. *American Journal of Human Genetics*, 60(3), 676–690.
- Backberg, M., Hervieu, G., Wilson, S., & Meister, B. (2002). Orexin receptor-1 (OX-R1) immunoreactivity in chemically identified neurons of the hypothalamus: Focus on orexin targets involved in control of food and water intake. *European Journal of Neuroscience*, 15(2), 315–328.
- Backman, D.R., Haddad, E.H., Lee, J.W., Johnston, P.K., & Hodgkin, G.E. (2002). Psychosocial predictors of healthful dietary behavior in adolescents. *Journal of Nutrition Educational Behavior*, 34(4), 184–192.
- Barberis, C., & Tribollet, E. (1996). Vasopressin and oxytocin receptors in the central nervous system. *Critical Reviews in Neurobiology*, 10(1), 119–154.
- Beck, B. (2000). Neuropeptides and obesity. *Nutrition*, 16(10), 916–923.
- Birch, L.L., McPhee, L., Sullivan, S., & Johnson, S. (1989). Conditioned meal initiation in young children. *Appetite*, 13(2), 105–113.
- Botta, R.A. (1999). Television images and adolescent girls' body image disturbance. *Journal of Communication*, 49(2), 22–41.
- Bulik, C.M., Sullivan, P.F., Wade, T.D., & Kendler, K.S. (2000). Twin studies of eating disorders: A review. *International Journal of Eating Disorders*, 27(1), 1–20.
- Clendenen, V.I., Herman, C.P., & Polivy, J. (1994). Social facilitation of eating among friends and strangers. *Appetite*, 23(1), 1–13.
- Coates, J.C., & de Bono, M. (2002). Antagonistic pathways in neurons exposed to body fluid regulate social feeding in *Caenorhabditis elegans*. *Nature*, 419(6910), 925–929.
- de Castro, J.M., & de Castro, E.S. (1989). Spontaneous meal patterns of humans: Influence of the presence of other people. *American Journal of Clinical Nutrition*, 50(2), 237–247.
- Devlin, B., Bacanu, S.A., Klump, K.L., Bulik, C.M., Fichter, M.M., Halmi, K.A., Kaplan, A.S., Strober, M., Treasure, J., Woodside, D.B., Berrettini, W.H., & Kaye, W.H. (2002). Linkage analysis of anorexia nervosa incorporating behavioral covariates. *Human Molecular Genetics*, 11(6), 689–696.
- Dhillon, W.S., Small, C.J., Jethwa, P.H., Russell, S.H., Gardiner, J.V., Bewick, G.A., et al. (2003). Paraventricular nucleus administration of calcitonin gene-related peptide inhibits food intake and stimulates the hypothalamo-pituitary-adrenal axis. *Endocrinology*, 144(4), 1420–1425.
- Frank, G.K., Kaye, W.H., Altemus, M., & Greeno, C.G. (2000). CSF oxytocin and vasopressin levels after recovery from bulimia nervosa and anorexia nervosa, bulimic subtype. *Biological Psychiatry*, 48(4), 315–318.
- Fulker, D.W., Cherny, S.S., Sham, P.C., & Hewitt, J.K. (1999). Combined linkage and association sib-pair analysis for quantitative traits. *American Journal of Human Genetics*, 64(1), 259–267.
- Garner, D.M. (1991). *Eating Disorders Inventory-2: Professional manual*. Odessa, FL: Psychological Assessment Resources.
- Garner, D.M., & Garfinkel, P.E. (1979). The Eating Attitudes Test: An index of the symptoms of anorexia nervosa. *Psychological Medicine*, 9(2), 273–279.
- Garner, D.M., & Garfinkel, P.E. (1980). Socio-cultural factors in the development of anorexia nervosa. *Psychological Medicine*, 10(4), 647–656.
- Garner, D.M., Olmsted, M.P., Bohr, Y., & Garfinkel, P.E. (1982). The Eating Attitudes Test: Psychometric features and clinical correlates. *Psychological Medicine*, 12, 871–878.

- Garner, D.M., Olmsted, M.P., & Garfinkel, P.E. (1985). Similarities among bulimic groups selected by different weights and weight histories. *Journal of Psychiatric Research*, 19(2-3), 129-134.
- Grice, D.E., Halmi, K.A., Fichter, M.M., Strober, M., Woodside, D.B., Treasure, J.T., et al. (2002). Evidence for a susceptibility gene for anorexia nervosa on chromosome 1. *American Journal of Human Genetics*, 70(3), 787-792.
- Hammock, E.A., & Young, L.J. (2002). Variation in the vasopressin V1a receptor promoter and expression: Implications for inter- and intraspecific variation in social behaviour. *European Journal of Neuroscience*, 16(3), 399-402.
- Hinney, A., Remschmidt, H., & Hebebrand, J. (2000). Candidate gene polymorphisms in eating disorders. *European Journal of Pharmacology*, 410, 147-159.
- Holland, A.J., Sicotte, N., & Treasure, J. (1988). Anorexia nervosa: Evidence for a genetic basis. *Journal of Psychosomatic Research*, 32(6), 561-571.
- Horeish, N., Apter, A., Ishai, J., Danziger, Y., Miculincer, M., Stein, D., et al. (1996). Abnormal psychosocial situations and eating disorders in adolescence. *Journal of the American Academy of Child and Adolescent Psychiatry*, 35(7), 921-927.
- Hsu, L.K. (1990). Experiential aspects of bulimia nervosa. Implications for cognitive behavioral therapy. *Behavior Modification*, 14(1), 50-65.
- Inenaga, K., Dyball, R.E., Okuya, S., & Yamashita, H. (1986). Characterization of hypothalamic noradrenaline receptors in the supraoptic nucleus and periventricular region of the paraventricular nucleus of mice in vitro. *Brain Research*, 369(1-2), 37-47.
- Inenaga, K., & Yamashita, H. (1986). Excitation of neurones in the rat paraventricular nucleus in vitro by vasopressin and oxytocin. *Journal of Physiology*, 370, 165-180.
- Insel, T.R., & Young, L.J. (2000). Neuropeptides and the evolution of social behavior. *Current Opinions in Neurobiology*, 10(6), 784-789.
- Kaye, W.H. (1996). Neuropeptide abnormalities in anorexia nervosa. *Psychiatry Research*, 62(1), 65-74.
- Kaye, W.H., Klump, K.L., Frank, G.K., & Strober, M. (2000). Anorexia and bulimia nervosa. *Annual Review in Medicine*, 51, 299-313.
- Kaye, W.H., & Strober, M. (1999). Neurobiology of eating disorders. In D.E. Charney, E.J. Nestler, & B.S. Bunney (Eds.), *Neurobiological foundations of mental illness* (pp. 891-906). New York: Oxford University Press.
- Kim, S.J., Young, L.J., Gonen, D., Veenstra-VanderWeele, J., Courchesne, R., Courchesne, E., et al. (2002). Transmission disequilibrium testing of arginine vasopressin receptor 1A (*AVPR1A*) polymorphisms in autism. *Molecular Psychiatry*, 7(5), 503-507.
- Klump, K.L., Miller, K.B., Keel, P.K., McGue, M., & Iacono, W.G. (2001). Genetic and environmental influences on anorexia nervosa syndromes in a population-based twin sample. *Psychological Medicine*, 31(4), 737-740.
- Kow, L.M., & Pfaff, D.W. (1986). Vasopressin excites ventromedial hypothalamic glucose-responsive neurons in vitro. *Physiology and Behaviour*, 37(1), 153-158.
- Lange, C., & Laird, N.M. (2002). Power calculations for a general class of family-based association tests: Dichotomous traits. *American Journal of Human Genetics*, 71(3), 575-584.
- Langhans, W., Delprete, E., & Scharrer, E. (1991). Mechanisms of vasopressin's anorectic effect. *Physiology and Behaviour*, 49(1), 169-176.
- Lesch, K.P., Bengel, D., Heils, A., Sabol, S.Z., Greenberg, B.D., Petri, S., et al. (1996). Association of anxiety-related traits with a polymorphism in the serotonin transporter gene regulatory region [see comments]. *Science*, 274(5292), 1527-1531.
- Levine, A.S., & Billington, C.J. (1997). Why do we eat? A neural systems approach. *Annual Review in Nutrition*, 17, 597-619.
- Levine, S. (2001). Primary social relationships influence the development of the hypothalamic-pituitary-adrenal axis in the rat. *Physiology and Behaviour*, 73(3), 255-260.
- Livesley, W.J., Jang, K.L., & Vernon, P.A. (1998). Phenotypic and genetic structure of traits delineating personality disorder [see comments]. *Archives of General Psychiatry*, 55(10), 941-948.
- Maes, M., Monteleone, P., Bencivenga, R., Goossens, F., Maj, M., van West, D., et al. (2001). Lower serum activity of prolyl endopeptidase in anorexia and bulimia nervosa. *Psychoneuroendocrinology*, 26(1), 17-26.
- Matsushita, S., Nakamura, T., Nishiguchi, N., & Higuchi, S. (2002). Association of serotonin transporter regulatory region polymorphism and abnormal eating behaviors. *Molecular Psychiatry*, 7(6), 538-540.
- Monge-Rojas, R., Nunez, H.P., Garita, C., & Chen-Mok, M. (2002). Psychosocial aspects of Costa Rican adolescents' eating and physical activity patterns. *Journal of Adolescent Health*, 31(2), 212-219.
- Monks, S.A., Kaplan, N.L., & Weir, B.S. (1998). A comparative study of sibship tests of linkage and/or association. *American Journal of Human Genetics*, 63(5), 1507-1516.
- Morimoto, I., Yamamoto, S., Kai, K., Fujihira, T., Morita, E., & Eto, S. (2000). Centrally administered murine-leptin stimulates the hypothalamus-pituitary-adrenal axis through arginine-vasopressin. *Neuroendocrinology*, 71(6), 366-374.
- Niimi, M., Murao, K., & Taminato, T. (2001). Central administration of neuromedin U activates neurons in ventrobasal hypothalamus and brainstem. *Endocrine*, 16(3), 201-206.
- Polivy, J., & Herman, C.P. (1987). Diagnosis and treatment of normal eating. *Journal of Consulting and Clinical Psychology*, 55(5), 635-644.
- Rabinowitz, D. (1997). A transmission disequilibrium test for quantitative trait loci. *Human Heredity*, 47(6), 342-350.

- Rutherford, J., McGuffin, P., Katz, R.J., & Murray, R.M. (1993). Genetic influences on eating attitudes in a normal female twin population. *Psychological Medicine*, 23(2), 425–436.
- Saper, C.B., Chou, T.C., & Elmquist, J.K. (2002). The need to feed: Homeostatic and hedonic control of eating. *Neuron*, 36(2), 199–211.
- Satia-Abouta, J., Patterson, R.E., Kristal, A.R., Teh, C., & Tu, S.P. (2002). Psychosocial predictors of diet and acculturation in Chinese American and Chinese Canadian women. *Ethnic Health*, 7(1), 21–39.
- Slatkin, M., & Excoffier, L. (1996). Testing for linkage disequilibrium in genotypic data using the expectation-maximization algorithm. *Heredity*, 76 (Pt 4), 377–383.
- Stice, E., Schupak-Neuberg, E., Shaw, H.E., & Stein, R.I. (1994). Relation of media exposure to eating disorder symptomatology: An examination of mediating mechanisms. *Journal of Abnormal Psychology*, 103, 836–840.
- Thibonnier, M., Graves, M.K., Wagner, M.S., Chatelain, N., Soubrier, F., Corvol, P., et al. (2000). Study of V(1)-vascular vasopressin receptor gene microsatellite polymorphisms in human essential hypertension. *Journal of Molecular Cell Cardiology*, 32(4), 557–564.
- Tozzi, F., Bergen, A.W., & Bulik, C.M. (2002). Candidate gene studies in eating disorders. *Psychopharmacology Bulletin*, 36(3), 60–90.
- Tribollet, E., Arsenijevic, Y., & Barberis, C. (1998). Vasopressin binding sites in the central nervous system: Distribution and regulation. *Progress in Brain Research*, 119, 45–55.
- Ur, E., Wilkinson, D.A., Morash, B.A., & Wilkinson, M. (2002). Leptin immunoreactivity is localized to neurons in rat brain. *Neuroendocrinology*, 75(4), 264–272.
- Van Ree, J.M. (1986). Role of pituitary and related neuropeptides in alcoholism and pharmacodependence. *Progress in Neuropsychopharmacology and Biological Psychiatry*, 10(2), 219–228.
- Walters, E.E., & Kendler, K.S. (1995). Anorexia nervosa and anorexic-like syndromes in a population-based female twin sample. *American Journal of Psychiatry*, 152(1), 64–71.
- Wang, Z., Young, L.J., De Vries, G.J., & Insel, T.R. (1998). Voles and vasopressin: A review of molecular, cellular, and behavioral studies of pair bonding and paternal behaviors. *Progress in Brain Research*, 119, 483–499.
- Welch, G.N., Upchurch, G.R., Jr., & Loscalzo, J. (1997). Homocysteine, oxidative stress, and vascular disease. *Hospital Practice (Off Ed)*, 32(6), 81–82, 85, 88–92.
- Wren, A.M., Small, C.J., Abbott, C.R., Jethwa, P.H., Kennedy, A.R., Murphy, K.G., et al. (2002). Hypothalamic actions of neuromedin U. *Endocrinology*, 143(11), 4227–4234.
- Yarkov, A., Montero, S., Lemus, M., Rocas de Alvarez-Buylla, E., & Alvarez-Buylla, R. (2001). Arginine-vasopressin in nucleus of the tractus solitarius induces hyperglycemia and brain glucose retention. *Brain Research*, 902(2), 212–222.